

## THE DISTRIBUTION AND ECOLOGICAL PREFERENCES OF STONE CRAYFISH *AUSTROPOTAMOBIOUS TORRENTIUM* (SCHRANK, 1803) (DECAPODA: ASTACIDAE) IN THE NORTH- WEST ROMANIA

LUCIAN PÂRVULESCU

**Abstract.** This paper presents data on geographical distribution and ecological preferences of *Austropotamobius torrentium* species following investigations of 104 sampling sites in the west and north-western Romania. Specimens were caught by actively searching the river bed, after which they were released. A detailed map of the distribution was prepared: the species was found to be living in 23 sampling sites from three basins: Crișul Alb, Crișul Negru and Crișul Repede. The current distribution of the species extends to the central and eastern part of the Codru Moma Mountains, western Bihor Mountains, eastern Pădurea Craiului Mountains, central and northern areas of the Vlădeasa Mountains and southern and central areas of the Plopiș Mountains. The northernmost population of Romania is found in the Valea Mare River, on the Plopiș Mountains. Generally, the species occurs on a substrate consisting primarily of stone and gravel. The species' preferred combination of riparian vegetation is *Alnus* sp. and *Salix* sp. in deciduous forests. An important number of investigated rivers, totalling 37, are occupied by the *Astacus astacus* species, data proving that there are no mixed populations. No other non-indigenous crayfish species were found during this study.

**Résumé.** Ce travail présente des données sur la distribution géographique et les préférences écologiques d'espèce *Austropotamobius torrentium* suite aux investigations sur 104 sites d'échantillonnage dans l'ouest et au nord-ouest de la Roumanie. Les spécimens ont été capturés par les chercheurs activement au fond de la rivière, après ils ont été libérés. Une carte détaillée de la distribution a été préparé: l'espèce a été trouvée dans 23 rivières de trois bassins: Crișul Alb, Crișul Negru et Crișul Repede. La répartition actuelle de l'espèce s'étend à la partie centrale et orientale des Montagnes Codru Moma, dans l'ouest des Montagnes Bihor, la partie orientale des Montagnes Pădurea Craiului, la zone centre et du nord de la Montagne Vlădeasa et le centre et la zone du sud des Montagnes Plopiș. L'espèce se trouve généralement dans les rivières avec un substrat constitué principalement de pierres de taille variable et de gravier. La combinaison de la végétation riveraine préférée par l'espèce étudiée est représenté par: *Alnus* sp. et *Salix* sp., dans les forêts à feuilles caduques. La plus nordique population d'espèce *A. torrentium* de la Roumanie se trouve dans la rivière Valea Mare, dans les Montagnes Plopiș. Dans un nombre important de rivières étudiées, totalisant 37, a été trouvée l'espèce *Astacus astacus*. Ces données montrent qu'il n'y a pas des populations mixtes. Dans cette étude, aucune des espèces envahissantes n'a pas été trouvé dans les zones étudiées.

**Key words:** *Astacus astacus*, *Austropotamobius torrentium*, crayfish, distribution, ecology, endangered species, noble crayfish, stone crayfish.

### INTRODUCTION

The stone crayfish *Austropotamobius torrentium* (Schrank, 1803) is one of the three indigenous crayfish species of the Romanian freshwater ecosystems, the others being the noble crayfish *Astacus astacus* (Linnaeus, 1758), and the narrow-clawed crayfish *Astacus leptodactylus* Eschscholtz, 1823 (Băcescu, 1967; Holdich et al., 2009; Pârvulescu, 2010). One non-native crayfish, the spiny-cheek crayfish *Orconectes limosus* (Rafinescque, 1817) (Pârvulescu et al., 2009), occurs in Romanian freshwater ecosystems. In the European Council's Directive 92/43 (assumed by Romanian law 57 / 2007), *A. torrentium* is rated as a "priority species",

and therefore it is necessary to implement a series of management measures in the protected areas, among which the periodic measurement of population density is of great importance (Pârvulescu, 2007). Now, the actual IUCN status rates this species as “data deficient” (Füreder et al., 2010).

The European distribution of this species covers the central and south-eastern areas of the continent, from Germany and the Czech Republic in the north, Luxembourg and France to the west, Greece to the south, and Romania, Bulgaria and Turkey in the east (Holdich et al., 2006). *Austropotamobius torrentium* usually lives in small and medium-sized rivers and streams (Vlach et al., 2009), requiring pristine waters (Svobodová et al., 2008; Pârvulescu et al., 2011). Over a period of decades, many factors could influence the geographical distribution of *A. torrentium*, for example, modification of its habitat and water pollution (Lowery & Hogger, 1986; Gherardi and Holdich, 1999; Füreder et al., 2003; Lyons & Kelly-Quinn, 2003), a lower rate of growth (Streissl & Hödl, 2002) and lower fecundity (Maguire et al., 2010). Anthropogenic impacts such as organic pollution and river bed modification might have triggered the disappearance of the species from sections of the watercourse situated downstream of villages and towns (Pârvulescu et al., 2011). The high pressure caused by both invasive crayfish species and the crayfish plague pathogen (*Aphanomyces astaci*) has become a serious threat, which is almost unstoppable based on present knowledge. At the end of 2011, *Orconectes limosus* already occupies the first 105 km of the Romanian Danube, carrying the crayfish plague (Pârvulescu et al., 2012).

There are few data published regarding the freshwater crayfish species in Romania. The earliest records of *A. torrentium* in Romania place this species in the south-west of the country (Scriban, 1908). Entz (1912) also recorded the species in the south-west area, and additionally in the centre of the Transylvanian Plateau. Between 1935 and 1962 there were references to its presence in the south-west, west and north-west of Romania (Băcescu, 1967). The catalogue of the “Grigore Antipa” National Museum of Natural History in Bucharest lists the localities where the species was found, mainly based on Mihai Băcescu’s collection (Petrescu & Petrescu, 2010). These earlier observations are difficult to georeference because geographical coordinates were not given, and so it is hard to establish a clearly defined geographical area for the distribution of *A. torrentium*. Also, as these records are over 30 years old, it is very possible that the actual situation may no longer be the same. This paper provides evidence for and completes the data relating to the distribution and ecology, in order to assist in efficient conservation management for this species.

#### MATERIAL AND METHODS

During 2010 and 2011, field sampling was carried out in order to establish the distribution of *A. torrentium* in the permanent waters in the upper sections of the main courses and tributaries from the mountain areas and sub-mountain areas of the west and north-west of Romania. The extent of the investigated area is from north of the Mureş River, covering 14 geographical units (i.e. mountains, plateaus, hills). Tributaries of six hydrographic basins were investigated: the Crişul Alb, Crişul Negru, Crişul Repede, Someş, Mureş and Tisa rivers. The sections for sampling sites were chosen at random, and we investigated an average of 15 mainstreams per geographical unit, with the number for each depending on its size. Each sampling site contained, on average, a stretch of river of approximately 200

m in length. The crayfish were collected using direct hand sampling from the river bed, by searching shores galleries, spaces between rocks and within the roots. The relative abundance (catch per unit effort, CPUE) of crayfish was estimated as the number of crayfish caught per 100 m length of river stretch. Where no crayfish were found, the sampling site was recorded as one with “no crayfish”, but this was only confirmed after investigating at least 300 m of a river. The captured crayfish specimens were released at exactly the same location at which they were caught. The riverbed morphology, shelter use and riparian vegetation were recorded when observations of the presence of crayfish were made. These indicators were defined in terms of the percentages of different types in each sector. The bottom types included boulders, stones, gravel, sand and mud, and the shelter use was defined accordingly to the type where each specimen was found.

A digital distribution map at 1:50,000 scale was prepared, based on the georeferenced field data. Using GlobalMapper software, we obtained a map of distribution points by marking where we found each of the crayfish species, as well as the sites where none were found. Then, using InkScape, we coloured the distribution points to make the areas visible. The most-probable distribution area was established by analysis of the distribution of the points, taking into consideration the biogeography of the species and the watershed area. Ecological parameters were studied with the weighted average method. The weights were represented by the relative abundance (CPUE).

### RESULTS AND DISCUSSION

In 60 of 104 investigated sampling sites, crayfish were present (Fig. 1), 23 populated by the stone crayfish (*Austropotamobius torrentium*) and 37 by the noble crayfish (*Astacus astacus*). Table 1 contains data summaries for the ecological parameters measured in the locations where *A. torrentium* specimens were found. The

Table 1

Data summary for several parameters of the habitat measured at the sampling sites where *A. torrentium* was found.  
Intervals with the highest frequencies are shaded.

River bottom mixture								
Type	boulders	stones	gravel	sand	mud			
Frequency (%)	6.8	31.8	43.5	12.7	5.2			
Shelter use								
Type	bank galleries	under stones		between roots	free			
Frequency (%)	28.6	54.2		16.8	0.4			
Tree coverage								
Interval	0-20 %	21-40 %	41-60 %	61-80 %	81-100 %			
Frequency (%)	11.2	18	10.1	25.9	34.8			
Riparian trees composition								
Genus or type	Coniferous or mixed	<i>Fagus</i> sp.	<i>Carpinus</i> sp.	<i>Quercus</i> sp.	<i>Corylus</i> sp.	<i>Alnus</i> sp.	<i>Salix</i> sp.	Other
Frequency (%)	2.9	5.5	6.8	0.3	6.2	37.4	24.5	16.4




Fig. 1 - The distribution area of *Austropotamobius torrentium* and *Astacus astacus* obtained from data assembled during the 2010-2011 fieldwork in the west and north-west of Romania.




Fig. 2 - The frequency of crayfish species in the hydrographic basins, investigated during the 2010-2011 fieldwork in the west and north-west of Romania.

species occurs on a substrate consisting primarily of stone and gravel, the preferred shelter is represented by the spaces under and between stones. The most frequent combination of riparian vegetation is *Alnus* sp. and *Salix* sp. These ecological results are similar to those presented in other publications (e.g. Machino & Füreder, 2005).

The distribution results are following described according to the hydrographic basin from which they were obtained, from south to north.

*The Mureş tributaries.* Twenty-five sampling sites of the middle section of the Mureş River were investigated from the Trascău, Metaliferi and Zărandului mountains indicating the presence of only *A. astacus* crayfish species (Tab. 2).

*The Crişul Alb hydrographic basin.* Twenty sampling sites distributed in the area of the Codru Moma, Bihor, Zărand and Metaliferi mountains were investigated; *A. torrentium* was found in 10% of the investigated sites, *A. astacus* was identified in 35% of the investigated sites, and 55% did not harbour crayfish (Fig. 2, tab. 2). The highest altitude at which *A. torrentium* was found was at 370 m, in the Valea Megheş River (Tab. 2).

*The Crişul Negru hydrographic basin.* Twenty-three sampling sites, situated in five geographical units, were investigated: the Vlădeasa, Plopiş, Pădurea Craiului, Bihor and Codru Moma mountains. Two crayfish species were identified: *A. torrentium* and *A. astacus*, the former being found in 47.8% of the investigated sites (Fig. 2, tab. 2). The location at the highest altitude was in the Boga Stream, at 560 m, where we captured only one specimen. *A. astacus* was captured in three sampling sites (Tab. 2).

*The Crişul Repede hydrographic basin.* Fourteen sampling sites in several geographical units in this basin were investigated: the Meseş, Plopiş, Pădurea Craiului and Vlădeasa mountains. *A. torrentium* was found in 71.4% of the investigated sites (Fig. 2, tab. 2). The maximum altitude at which the species was found was 800 m, in the Valea Ieduşului Stream. The highest population density registered for the species in the investigated area was 55.8 individuals per 100 m of river stretch on the Valea Mare River. *A. astacus* was captured in four sampling sites (Tab. 2).

*The Someş hydrographic basin.* Fourteen sampling sites were investigated in the area of the Gutâi, Țibleş, Meseş, and Gilău mountains and western part of the Transylvanian Plateau. Only *A. astacus* was found (Tab. 2).

*The Tisa tributaries.* Eight sampling sites were investigated, situated on the Oaş, Gutâi and Țibleş mountains, only *A. astacus* being found (Tab. 2).

Compared to populations of *A. torrentium* from the south-west of Romania (Pârvulescu & Petrescu, 2010), the western and north-western populations are scarcely represented. These populations occupy the central and eastern part of the Codru Moma Mountains, the western Bihor Mountains, the eastern Pădurea Craiului Mountains, the central and northern Vlădeasa Mountains and the southern and central Plopiş Mountains (Fig. 2). The majority of this area is included in the Apuseni Nature Park and several Natura 2000 sites. The northern population of this species in Romania is represented by the Valea Mare River, in the Plopiş Mountains. Contrary to information presented by the collection of “Grigore Antipa” National Museum of Natural History of Bucharest (Petrescu & Petrescu, 2010), the species was not found in the Tisa tributaries, where only *A. astacus* was present.

The non-indigenous *O. limosus* does not represent a significant threat for *A. torrentium* in the west and north-west of Romania. It is known that this invasive species spreads through large rivers, and colonization of small tributaries is very rare, and only through human mediation (Petrušek et al., 2006; Pârvulescu et al., 2012). The pathway for *O. limosus* colonization in the future could be the Mureş and Tisa

Table 2

Crayfish sampled rivers toponyms, geographic coordinates (Stereo 70), and captured specimens.

Rivers toponyms	Geographic coordinates	Altitude (m)	<i>Austropotamobius torrentium</i>	<i>Astacus astacus</i>
<b>Mureș tributaries</b>				
Tămășești	46°01'32"N / 22°30'39"E	340	-	1 ♂, 5 ♀♀
Carpăn	46°03'58"N / 22°55'51"E	179	-	2 ♂♂, 1 ♀
Căianu	46°02'00"N / 22°53'05"E	276	-	4 ♂♂, 5 ♀♀
Băcaia	46°00'59"N / 23°10'27"E	370	-	2 ♂♂, 6 ♀♀
Conop	46°07'34"N / 21°53'13"E	180	-	2 ♂♂
Cladovița	46°09'07"N / 21°41'07"E	200	-	6 ♀♀
Balșa	46°03'12"N / 23°04'32"E	360	-	2 ♂♂, 6 ♀♀
Geoagiu	46°01'12"N / 23°07'17"E	330	-	9 ♂♂, 11 ♀♀
Valea Țâlnei	46°10'47"N / 22°29'16"E	420	-	3 ♂♂
Văleni	46°05'22"N / 23°25'37"E	340	-	1 ♀
Văltorii	46°08'29"N / 23°12'54"E	690	-	2 ♂♂, 1 ♀
<b>Crișul Alb hydrographic basin</b>				
Valea Stoiaca	46°12'59"N / 21°44'23"E	250	-	1 ♂
Nadăș	46°12'38"N / 21°55'20"E	240	-	4 ♂♂, 10 ♀♀
Valea Megheș	46°28'14"N / 22°15'55"E	370	1 ♂, 1 ♀	-
Țebea	46°08'46"N / 22°42'08"E	350	-	3 ♂♂, 4 ♀♀
Valea Birtinului	46°09'40"N / 22°38'40"E	290	-	3 ♀♀
Derjana	46°13'49"N / 22°42'32"E	280	-	2 ♀♀
Rănușa	46°26'21"N / 22°16'02"E	250	2 ♂♂, 3 ♀♀	-
Valea Prundu	46°12'37"N / 22°06'16"E	340	-	1 ♀
Aciua	46°21'46"N / 22°29'00"E	270	-	3 ♂♂, 4 ♀♀
<b>Crișul Negru hydrographic basin</b>				
Valea Archișel	46°30'22"N / 22°07'53"E	390	-	3 ♀♀
Valea Zerzegului	46°37'47"N / 22°09'00"E	450	1 ♂, 2 ♀♀	-
Valea Armanului	46°38'43"N / 22°13'38"E	300	1 ♂	-
Sohodol	46°50'41"N / 22°22'21"E	350	1 ♀, 3 ♀♀	-
Cuților	46°51'14"N / 22°25'17"E	525	2 ♀♀	-
Tâlniciorii	46°25'00"N / 22°27'48"E	480	3 ♂♂, 6 ♀♀	-
Valea Rudăreasa	46°28'10"N / 22°31'00"E	360	2 ♂♂, 2 ♀♀	-
Valea Sighiștel	46°31'33"N / 22°33'32"E	380	1 ♀	-
Racăș	46°51'51"N / 22°18'42"E	220	-	1 ♂, 2 ♀♀

Table 2 (continued)

Rivers toponyms	Geographic coordinates	Altitude (m)	<i>Austropotamobius torrentium</i>	<i>Astacus astacus</i>
Crăiasa	46°32'48"N / 22°35'41"E	530	5 ♂♂, 4 ♀♀	-
Boga	46°36'31"N / 22°39'33"E	560	1 ♀	-
Valea Finişului	46°34'10"N / 22°13'50"E	445	1 ♀	-
Valea Mare	46°38'30"N / 22°14'41"E	250	-	2 ♂♂, 2 ♀♀
Şoimuşurilor	46°48'31"N / 22°25'27"E	350	3 ♂♂	-
<b>Crişul Repede hydrographic basin</b>				
Luncşoru	47°06'27"N / 22°36'44"E	380	4 ♂♂, 3 ♀♀	-
Valea Mare	47°07'39"N / 22°36'39"E	310	26 ♂♂, 28 ♀♀	-
Poicu	46°59'05"N / 22°55'12"E	625	3 ♂♂, 3 ♀♀	-
Valea Boului	47°01'52"N / 22°51'27"E	410	-	1 ♀
Barcău	47°03'01"N / 22°44'50"E	425	-	3 ♀♀
Valea Peştilor	47°06'11"N / 22°39'39"E	315	-	3 ♂♂, 1 ♀
Valea Ieduţului	46°41'28"N / 22°35'11"E	800	4 ♂♂, 5 ♀♀	-
Valea Runcu	46°45'40"N / 22°34'03"E	630	2 ♂♂, 1 ♀	-
Răchiteasca	47°03'30"N / 22°31'51"E	290	1 ♂, 3 ♀♀	-
Valea Gârbului	46°56'01"N / 22°52'21"E	460	1 ♀	-
Valea Vişagului	46°51'51"N / 22°48'52"E	590	1 ♀	-
Valea Bisericii	46°50'08"N / 22°40'13"E	495	2 ♂♂, 2 ♀♀	-
Damiş	46°52'00"N / 22°32'12"E	700	2 ♂♂, 4 ♀♀	-
Valea Mierei	46°58'05"N / 22°24'30"E	455	-	1 ♂, 1 ♀
<b>Someş hydrographic basin</b>				
Vicleanul Mare	47°40'29"N / 23°36'26"E	290	-	3 ♂♂, 1 ♀
Valea Usturoiului	47°41'04"N / 23°34'24"E	320	-	3 ♀♀
Valea Mare	47°43'06"N / 23°25'05"E	280	-	1 ♂
Agrij	47°01'10"N / 22°58'51"E	425	-	1 ♂, 2 ♀♀
Gârbău	46°43'54"N / 23°31'52"E	350	-	1 ♂
Someşul Cald	46°43'50"N / 23°18'47"E	430	-	3 ♀♀
Tetişu	46°54'45"N / 23°03'22"E	340	-	1 ♂
<b>Tisa tributaries</b>				
Valea Holia	48°05'07"N / 23°13'03"E	210	-	3 ♀♀
Valea Mare	48°00'46"N / 23°18'21"E	200	-	1 ♂
Sărăsău	47°56'53"N / 23°47'52"E	300	-	5 ♂♂, 7 ♀♀
Băleaşa	47°38'14"N / 24°20'59"E	540	-	1 ♀
Valea Tejei	47°48'03"N / 23°53'29"E	440	-	1 ♂, 1 ♀

rivers. In the Hungarian lower course of the Tisa River, the presence of *O. limosus* has already been reported (Puky & Schád, 2006; Puky, 2009). The worst threat, represented by *Pacifastacus leniusculus*, should be taken into consideration over the next decades. In Hungary, this species is reported in Rába River (Puky et al., 2005), and it is known to be more susceptible to colonization of the smaller tributaries, even brooks and springs (Holdich et al., 2006): this creates the possibility of contact with *A. torrentium*, and therefore transmission of the crayfish plague infection to their populations, considerably higher.

#### ACKNOWLEDGEMENTS

This study was funded by CNCIS - Exploratory research projects PCE-4 grant no 1458/2008 "The stone crayfish (*Austropotamobius torrentium*), distribution in Romanian habitats, ecology and genetics of populations". I want to thank the Romanian Academy, especially The Commission for Natural Protected Monuments, for approving this study, as well as the administrators of the natural protected areas for permission to work in the field. Also, many thanks to Zoltan Kovacs and Zoltan Mate for their help in the Plopiş Mountains investigations. Last, but not least, I want to thank to the volunteer students of the Faculty of Chemistry, Biology and Geography from the West University of Timișoara for participation in the field activities. The anonymous reviewers provided useful suggestions on a previous version of the manuscript.

#### DISTRIBUȚIA ȘI PREFERINȚELE ECOLOGICE ALE RACULUI-DE-PONOARE *AUSTROPOTAMOBIUS TORRENTIUM* (SCHRANK, 1803) (DECAPODA: ASTACIDAE) ÎN NORD-VESTUL ROMÂNIEI

#### REZUMAT

În această lucrare sunt prezentate date referitoare la distribuția geografică și preferințele ecologice ale speciei *Austropotamobius torrentium* în urma investigării a 104 stații de prelevare din zona de vest și de nord-vest a României. Exemplarele au fost capturate activ, căutând în albia râului, după care au fost eliberate. A fost realizată o hartă detaliată de distribuție, specia fiind găsită la 23 din stațiile de prelevare făcând parte din bazinele râurilor Crișul Alb, Crișul Negru și Crișul Repede. Aria actuală de răspândire a speciei ocupă centrul și estul Munților Codru Moma, vestul Munților Bihor, estul Munților Pădurea Craiului, centrul și nordul Munților Vlădeasa, și sudul și centrul Munților Plopiș. Cea mai nordică populație din România este reprezentată de râul Valea Mare din Munții Plopiș. În general, specia ocupă cursuri de apă cu un substrat format din pietre și pietriș. Combinația de vegetație ripariană preferată este *Alnus* sp. și *Salix* sp. în etajul pădurilor de foioase. La un număr important de pâraie investigate, 37 din numărul total, a fost găsită specia *Astacus astacus*, datele arătând că nu există populații mixte. Nicio specie invazivă nu a fost găsită în timpul acestui studiu.

#### LITERATURE CITED

- BĂCESCU, M. C., 1967 - Crustacea, Decapoda. *In*: Fauna Republicii Socialiste România. Edit. Academiei Republicii Socialiste România, București, 4: 209-219. (in Romanian)
- ENTZ, G., 1912 - Über die Flußkrebse Ungarns. *Mathematischen Naturwissenschaftlichen Berichte aus Ungarn*, 30 (2): 67-125.
- FÜREDER, L., F. GHERARDI, C. SOUTY-GROSSET, 2010 - *Austropotamobius torrentium*. *In*: IUCN 2011. IUCN Red List of Threatened Species. Version 2011.1. <www.iucnredlist.org>. Downloaded on 24 October 2011.
- FÜREDER, L., B. OBERKOFER, R. HANEL, J. LEITER, B. THALER, 2003 - The freshwater crayfish *Austropotamobius pallipes* in South Tyrol: heritage species and bioindicator. *Bulletin Français de la Pêche et de la Pisciculture*, 370-371: 79-95.
- GHERARDI, F., D. M. HOLDICH, 1999 - Crayfish in Europe as alien species. How to make the best of a bad situation. *Crustacean Issues* 11, A. A. Balkema, Rotterdam, Brookfield, 299 pp.
- HOLDICH, D. M., P. HAFFNER, P. Y. NOËL, 2006 - Species files. *In*: C. Souty-Grosset, D. M. Holdich, P. Y. Noël, J. D. Reynolds, P. Haffner (eds), Atlas of Crayfish in Europe. Muséum National d'Histoire Naturelle, Paris, (Patrimoines Naturels, 64): 49-129.
- HOLDICH, D. M., J. D. REYNOLDS, C. SOUTY-GROSSET, P. J. SIBLEY, 2009 - A review of the ever increasing threat to European crayfish from non-indigenous crayfish species. *Knowledge and Management of Aquatic Ecosystems*, 394-395, 11: 1-46.

- LOWERY, R., J. HOGGER, 1986 - The effect of river engineering and disease on a population of *Austropotamobius pallipes* in the River Lea, UK. *Freshwater Crayfish*, 6: 94-97.
- LYONS, R., M. KELLY-QUINN, 2003 - An investigation into the disappearance of *Austropotamobius pallipes* (Lereboullet) populations in the headwaters of the Nore River, Ireland and the correlation to water quality. *Bulletin Français de la Pêche et de la Pisciculture*, 370-371: 139-150.
- MACHINO, Y., L. FÜREDER, 2005 - How to find a stone crayfish *Austropotamobius torrentium* (Schränk, 1803): a biogeography study in Europe. *Bulletin Français de la Pêche et de la Pisciculture*, 376-377: 507-517.
- MAGUIRE, I., A. LUCIĆ, M. SCHLETTERER, D. SINT, R. ERBEN, L. FÜREDER, 2010 - Fecundity of Indigenous Stone Crayfish, *Austropotamobius torrentium*, Populations: Implications for Crayfish Conservation in Austria and Croatia. *Freshwater Crayfish*, 17: 103-107.
- PÂRVULESCU, L., 2007 - *Austropotamobius torrentium*. Pp. 47-49. In: I. Combroux, E. Thiry, T. Toia (eds), *Caie de habitate și specii - Fișe pilot*. Edit. Balcanic, Timișoara. (in Romanian)
- PÂRVULESCU, L., 2010 - Crayfish field guide of Romania. Edit. Bioflux, Cluj-Napoca: 28 p.
- PÂRVULESCU, L., I. PETRESCU, 2010 - The distribution of stone crayfish *Austropotamobius torrentium* (Schränk, 1803) (Crustacea: Decapoda: Astacidae) in the south-west Romanian mountain and sub-mountain area. *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"*, 53: 103-113.
- PÂRVULESCU, L., O. PACIOGLU, C. HAMCHEVICI, 2011 - The assessment of the habitat and water quality requirements of the stone crayfish (*Austropotamobius torrentium*) and noble crayfish (*Astacus astacus*) species in the rivers from the Anina Mountains (SW Romania). *Knowledge and Management of Aquatic Ecosystems*, 401, 03: 1-12.
- PÂRVULESCU, L., C. PALOȘ, P. MOLNAR, 2009 - First record of spiny-cheek crayfish *Orconectes limosus* (Rafinesque, 1817) (Crustacea: Decapoda: Cambaridae). *North-Western Journal of Zoology*, 5 (2): 424-428.
- PÂRVULESCU, L., A. SCHRIMPF, E. KOZUBÍKOVÁ, S. C. RESINO, T. VRÁLSTAD, A. PETRUSEK, R. SCHULZ, 2012 - Invasive crayfish and crayfish plague on the move: first detection of the plague agent *Aphanomyces astaci* in the Romanian Danube. *Diseases of Aquatic Organisms*, 98: 85-94.
- PETRESCU, I., A. M. PETRESCU, 2010 - The catalogue of the freshwater crayfish (Crustacea: Decapoda: Astacidae) from Romania preserved in "Grigore Antipa" National Museum of Natural History of Bucharest. *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"*, 53: 115-123.
- PETRUSEK, A., L. FILIPOVA, Z. ĎURIS, I. HORKA, P. KOZAK, T. POLICAR, M. ŠTAMBERGOVA, Z. KUCERA, 2006 - Distribution of the invasive spiny-cheek crayfish (*Orconectes limosus*) in the Czech Republic. Past and present. *Bulletin Français de la Pêche et de la Pisciculture*, 380-381: 903-917.
- PUKY, M., 2009 - Confirmation of the presence of the spiny-cheek crayfish *Orconectes limosus* (Rafinesque, 1817) (Crustacea: Decapoda: Cambaridae) in Slovakia. *North-Western Journal of Zoology*, 5: 214-217.
- PUKY, M., P. SCHAD, 2006 - *Orconectes limosus* colonises new areas fast along the Danube in Hungary. *Bulletin Français de la Pêche et de la Pisciculture*, 380-381: 919-926.
- PUKY, M., J. D. REYNOLDS, P. SCHÁD, 2005 - Native and alien decapoda species in Hungary: distribution, status, conservation importance. *Bulletin Français de la Pêche et de la Pisciculture*, 376-377: 553-568.
- SCRIBAN, I. A., 1908 - Note sur les Ecrevisses de Roumanie. *Annales Scientifiques de l'Université de Jassy*, 5: 80-83.
- STREISSL, F., W. HÖDL, 2002 - Growth, morphometrics, size at maturity, sexual dimorphism and condition index of *Austropotamobius torrentium* Schränk. *Hydrobiologia*, 477: 201-208.
- SVOBODOVÁ, J., M. ŠTAMBERGOVÁ, P. VLACH, J. PICEK, K. DOUDA, M. BERÁNKOVÁ, 2008 - The impact of the water quality on the crayfish population in the Czech Republic, comparison with legislation of the Czech Republic. *VTEI*, 50: 1-5.
- VLACH, P., D. FISCHER, L. HULEC, 2009 - Microhabitat preferences of the stone crayfish *Austropotamobius torrentium* (Schränk, 1803). *Knowledge and Management of Aquatic Ecosystems*, 394-395, 15: 1-13.
- \*\*\* EUROPEAN COMMUNITIES, 1992 - Council Directive (92/43/EEC) on the conservation of natural habitats and of wild fauna and flora, The Council of European Communities.

*Received: October 27, 2011*  
*Accepted: July 23, 2012*

*West University of Timișoara*  
*Faculty of Chemistry, Biology, Geography*  
*Department of Biology-Chemistry*  
*Pestalozzi 16A St., 300115 Timișoara, Romania*  
e-mail: parvulescubio@cbg.uvt.ro